

How to separate chromosomes and restriction fragments using pulsed field gel electrophoresis.

David D. Perkins

Background

DNA content of the seven individual *Neurospora crassa* chromosomes ranges from 4 to 10.3 megabases (Orbach 1992). Because of chromosome size, the separation of intact whole-chromosome DNA by pulsed-field gel electrophoresis (PFGE) is slower and more difficult with *Neurospora* than with yeast (Volrath and Davis 1987) or other fungi that have much smaller chromosomes. Nevertheless, DNAs from the individual *Neurospora* chromosomes can be resolved successfully. Intact DNA as large as 14 megabases has been separated from one chromosome-rearrangement strain (Barry and Pollard 1993).

PFGE has been used in *Neurospora* to obtain estimates of chromosome and genome size (Orbach *et al.* 1988, Orbach 1992), to map translocation breakpoints (Smith and Glass 1996), and to isolate DNA from individual chromosomes (Ballario *et al.* 1989, Kelkar *et al.* 2001), thus enabling the generation of physical-clone maps of individual linkage groups by hybridization-based mapping (Aign *et al.* 2001). PFGE has also been used to separate restriction fragments in studies of chromosome breakage and of ribosomal DNA copy-number changes in the nucleolus organizer region (Butler 1991, 1992; Butler and Metzberg 1990, 1993).

Procedures

Based on earlier electrophoretic karyotyping methodology of Orbach *et al.* (1988), Kelkar *et al.* (2001) describe the equipment, materials, and conditions for the preparation, stepwise separation, and isolation of intact chromosomal DNAs representing the seven linkage groups of *N. crassa*. In wild type, the two largest DNAs (linkage groups I and V) and the two smallest (VI and VII) are too close in size to be well resolved. Separation of these DNAs was therefore accomplished using appropriate translocation strains.

The restriction fragments used by Butler and Metzberg in their studies were small enough to allow them to follow the protocol used for yeast by Chu *et al.* (1986) and Vollrath and Davis (1987). Electrophoresis was done for 24 hours in 0.5× TBE at 10–11°, with agarose gels cast in 0.5× TBE. Pulse times and voltages were varied to optimize resolution of the pertinent fragments.

References

Aign, V., U. Schulte, and J. D. Hoheisel. 2001. Hybridization-based mapping of *Neurospora crassa* linkage groups II and V. *Genetics* 157: 1015-1020.

Ballario, P., G. Morelli, E. Sporeno, and G. Macino. 1989. Cosmids from the Vollmer-Yanofsky library identified with a chromosome VII probe. *Fungal Genet. Newslett.* 36: 38-39.

Barry, E. G., and V. C. Pollard. 1993. Identification of *Neurospora* rearrangement chromosomes by PFGE. Proc. 17th Intern. Cong. Genet. (Moscow) Volume of Abstracts, p. 134.

Butler, D. 1991. Chromosome breakage and new telomere formation in the ribosomal DNA of *Neurospora crassa*. Fungal Genet. Newslett. 38: 49. (Abstr.).

Butler, D. K. 1992. Ribosomal DNA is a site of chromosome breakage in aneuploid strains of *Neurospora*. Genetics 131: 581-592.

Butler, D. K., and R. L. Metzenberg. 1990. Expansion and contraction of the nucleous organizer region of *Neurospora*: Changes originate in both proximal and distal segments. Genetics 126: 325-333.

Butler, D. K., and R. L. Metzenberg. 1993. Amplification of the nucleolus organizer region during the sexual phase of *Neurospora crassa*. Chromosoma 102: 519-525.

Chu, G., D. Voolrath, and R. W. Davis. 1986. Separation of large DNA molecules by contour-clamped homogeneous electric fields. Science 234: 1582-1584.

Kelkar, H. S., J. Griffith, M. E. Case, S. F. Covert, R. D. Hall, C. H. Keith, J. S. Oliver, M. J. Orbach, M. S. Sachs, J. R. Wagner, M. J. Weise, J. K. Wunderlich, and J. Arnold. The *Neurospora crassa* genome: Cosmid libraries sorted by chromosome. 2001. Genetics 157: 979-990.

Orbach, M. J. 1992. [Untitled note giving revised size estimates of *Neurospora* chromosomes.] Fungal Genet. Newslett. 39: 92.

Orbach, M. J., D. Vollrath, R. W. Davis, and C. Yanofsky. 1988. An electrophoretic karyotype of *Neurospora crassa*. Mol. Cell. Biol. 8: 1469-1473.

Smith, M. L., and N. L. Glass. 1996. Mapping translocation breakpoints by orthogonal field agarose-gel electrophoresis. Curr. Genet. 29: 301-305.

Vollrath, D., and R. W. Davis. 1987. Resolution of greater than 5 mega-basepair DNA molecules by contour-clamped homogeneous electric fields. Nucleic Acids Res. 15:7865-7876.

DDP